

ZERO CYCLES ON GENERIC HYPERSURFACES OF LARGE DEGREE

NAJMUDDIN FAKHRUDDIN

ABSTRACT. We show that given a smooth projective variety X over $\mathbb{C}$ with $\dim(X) \geq 3$, an ample line bundle $\mathcal{O}(1)$ on X and an integer $n > 1$, any n distinct points on a generic hypersurface of degree d are linearly independent in $CH_0(X)$ if $d \gg 0$. This generalizes a result of C. Voisin.

Let X be a smooth projective algebraic variety over $\mathbb{C}$ of dimension $r + 1 \geq 3$. Let $\mathcal{L}$ be an ample line bundle on X . Let $S^d = H^0(X, \mathcal{L}^d)$ and for a point $x \in X$, let $S_x^d = H^0(X, \mathcal{L}^d \otimes \mathcal{I}_x)$, where $\mathcal{I}_x$ is the ideal sheaf of x . Let $R = \bigoplus_{d \geq 0} S^d$.

Lemma 1. *There exists an integer $m > 0$ such that, for all $d \gg 0$, the following holds:*

- (i) *The natural map $S^d \otimes S^m \rightarrow S^{d+m}$ is surjective.*
- (ii) *The natural map $S^d \otimes S_x^m \rightarrow S_x^{d+m}$ is surjective.*

Proof. Let the ring R be generated in degrees $\leq t$. Let $m_1 = t!$. It is easy to see that $a \cdot m_1$ satisfies condition (i) of the lemma for $d > a \cdot t! \cdot t(t+1)/2$, for any positive integer a . Let d' be a positive integer so that $\mathcal{L}^{d'}$ is generated by its global sections. There exists m , a positive multiple of m_1 , such that the maps $S^{d'} \otimes S_x^m \rightarrow S_x^{d'+m}$ are surjective for all $x \in X$. For $d \gg 0$, both the maps $S^{d-d'} \otimes S^{d'} \rightarrow S^d$ and $S^{d-d'} \otimes S_x^{d'+m} \rightarrow S_x^{d+m}$ are surjective since $\mathcal{L}$ is ample, hence the map $S^{d-d'} \otimes S^{d'} \otimes S_x^m \rightarrow S_x^{d+m}$ is also surjective. The following commutative diagram then shows that the map $S^d \otimes S_x^m \rightarrow S_x^{d+m}$ is surjective.

$$\begin{array}{ccc} S^{d-d'} \otimes S^{d'} \otimes S_x^m & \longrightarrow & S^{d-d'} \otimes S_x^{d'+m} \\ \downarrow & & \downarrow \\ S^d \otimes S_x^m & \longrightarrow & S_x^{d+m} \end{array}$$

□

Let $\mathcal{X} \subset X \times S^d$ be the universal hyperplane section. For $s \in S^d$ we denote the fibre $p_2^*(s)$ by X_s , which we shall assume to be smooth. For a vector bundle $\mathcal{V}$ on X , by $\mathcal{V}(b)$ we shall mean $\mathcal{V} \otimes \mathcal{L}^b$.

Proposition 1. *For $d \gg 0$, the bundle $T\mathcal{X}(m)|_{X_s}$ is generated by its global sections.*

Proof. The proof, given the previous lemma, is identical to Proposition 1.1 of [1] and is hence omitted. □

Corollary 1. *There exists a linear function $d(n)$ of n , such that for all $d \geq d(n)$, the vector bundle $\Omega_{\mathcal{X}}^{dim S^d}|_{X_s}$ separates any n distinct points of X_s i.e. the global*

sections of the bundle surject onto the global sections of the bundle restricted to any subscheme consisting of n distinct reduced points.

Proof. $\Omega_{\mathcal{X}}^{dim S_d}|_{X_s} \cong \Omega_{\mathcal{X}}^r|_{X_s}^* \otimes \mathcal{K}_{\mathcal{X}}|_{X_s} \cong \wedge^r T\mathcal{X}|_{X_s} \otimes \mathcal{K}_{X_s} \cong \wedge^r T\mathcal{X}(r \cdot m)|_{X_s} \otimes \mathcal{K}_{X_s}(-r \cdot m) \cong \wedge^r T\mathcal{X}(r \cdot m)|_{X_s} \otimes \mathcal{K}_X(d - r \cdot m)|_{X_s}$. By the proposition $\wedge^r T\mathcal{X}(r \cdot m)|_{X_s}$ is generated by global sections if $d \gg 0$. Since $\mathcal{L}$ is ample, there exists a linear function $d(n)$ such that $\mathcal{K}_X(d - r \cdot m)|_{X_s}$ separates n points if $d \geq d(n)$. It follows that the tensor product also separates n distinct points. $\square$

Theorem 1. *Let X be a smooth projective variety of dimension $r + 1 \geq 3$ and let $\mathcal{L}$ be an ample line bundle on X . Then there exists a linear function $d(n)$ such that for all $d \geq d(n)$, any n distinct points of a generic hypersurface X_s , $s \in S^d$, are linearly independent in $CH^r(X_s)$.*

Proof. Suppose not. Then there exists an etale map $S \rightarrow S^d$ and n distinct sections $\sigma_1, \dots, \sigma_n$ of $\mathcal{X}_S$ such that the classes of these sections in $CH^r(\mathcal{X}_S)$ are linearly dependent. We may assume that S is affine and that all the fibres are smooth. Consider the classes of these cycles, $[\sigma_i]$, in the Hodge cohomology group $H^r(\mathcal{X}_S, \Omega_{\mathcal{X}_S}^r)$. By the Grothendieck-Serre duality, it is easy to see that as an element of $Hom(H^0(\mathcal{X}_S, \Omega_{\mathcal{X}_S}^{dim S}), H^0(S, \Omega_S^{dim S}))$, $[\sigma_i]$ is nothing but the restriction map σ_i^* on differential forms. By the previous corollary we see that all the σ_i^* are linearly independent, which is a contradiction. $\square$

Corollary 2. *Let X , $\mathcal{L}$ be as above. There exists a linear function $d'(n)$ such that for all $d \geq d'(n)$, the generic hypersurface X_s , $s \in S^d$, does not contain any (possibly singular) n -gonal curves.*

Proof. Follows easily from the theorem by considering two distinct elements in the linear system corresponding to a degree n map from the normalisation of the curve to $\mathbb{P}^1$. $\square$

Acknowledgements. This note was written while the author was partially supported by an N.S.F. grant at the Institute for Advanced Study, Princeton, during the year 1995-1996.

REFERENCES

- [1] C. VOISIN, *On a conjecture of Clemens on rational curves on hypersurfaces*, J. Differential Geom., 44 (1996), pp. 200–213.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

E-mail address: naf@math.tifr.res.in